

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau




(43) International Publication Date
26 July 2001 (26.07.2001)

PCT

(10) International Publication Number
WO 01/53340 A2

- (51) International Patent Classification: C07K 14/435 (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (21) International Application Number: PCT/US01/01740
- (22) International Filing Date: 19 January 2001 (19.01.2001)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data: 60/177,319 21 January 2000 (21.01.2000) US
- (71) Applicant (for all designated States except US): PHARMACIA & UPJOHN COMPANY [US/US]; 301 Henrietta Street, Kalamazoo, MI 49001 (US).
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- Published:
— without international search report and to be republished upon receipt of that report
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): GURNEY, Mark, E. [US/IS]; Tjarnagata 39, Reykjavik (IS). ABRAHAM, Irene [US/US]; 2576 Ninth Avenue West, Seattle, WA 98119 (US).
- (74) Agents: DELUCA, Mark et al.; Woodcock Washburn Kurtz Mackiewicz & Norris LLP, 46th Floor, One Liberty Place, Philadelphia, PA 19103 (US).
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: TRANSGENIC MOUSE MODEL OF HUMAN NEURODEGENERATIVE DISEASE



(57) Abstract: This invention provides transgenic mice expressing wild-type or mutant isoforms of human tau protein. The transgenic mice of the invention have many uses, including screening candidate drug compounds effective against neurodegenerative diseases and other diseases involving tauopathies. The transgenic mice of the invention can also be used in genetic crosses with other transgenic mice strains to model the progression of neurodegenerative disease.

WO 01/53340 A2

TRANSGENIC MOUSE MODEL OF HUMAN NEURODEGENERATIVE DISEASE

CROSS-REFERENCE TO RELATED APPLICATIONS

5 This application claims priority to U.S. Provisional Patent Application, Serial
Number 60/177,319 filed January 21, 2000, which is hereby incorporated by reference.

FIELD OF THE INVENTION

10 This invention is related to the fields of transgenic animals, neurodegenerative
disease, and of identifying compounds to treat such diseases.

BACKGROUND OF THE INVENTION

Many neurological diseases such as Alzheimer's disease, frontotemporal dementia
and Parkinsonism linked to chromosome 17 (FTDP-17), progressive supranuclear palsy,
15 corticobasal degeneration, and other undefined dementias, have the common pathological
feature of neurofibrillary tangles (NFTs). NFTs are structures made up of paired helical
filaments (PHFs), composed primarily of tau protein in a hyperphosphorylated state. Tau
belongs to a family of microtubule associated proteins (MAPs) that are thought to play
critical roles in stabilizing and remodeling the neuronal cytoskeleton. Under normal
20 physiological conditions tau is found primarily in the brain, associated with the cellular
cytoskeleton.

The adult human brain expresses six different isoforms of tau protein, ranging in
size from 352 to 441 amino acid residues. The isoforms differ by having either three or
four microtubule binding domains, and by the presence or absence of amino-terminal
25 inserts. Tau is phosphorylated in the brain by a variety of kinases, although the
relationship between hyperphosphorylation and tangle formation is unclear. It is also

unclear how plaque and tangle formation leads to cognitive dysfunction and neuronal loss in patients with Alzheimer's disease.

Insight into the relationship of tau pathology and dementia has come through the study of FTDP-17, which is caused by mutations in the *tau* gene (see generally Goedert *et al.*, 1998, *Neuron* 21:955-958). The signature neuropathy associated with FTDP-17 is insoluble filamentous aggregates of hyperphosphorylated tau protein occurring in the absence of amyloid plaque and Lewy bodies, hallmarks of Alzheimer's disease and Pick's disease, respectively. Tau deposits are present not only in neurons, but also in large numbers of glial cells including predominantly oligodendrocytes, and also astrocytes (Spillantini *et al.*, 1997, *Proc. Natl. Acad. Sci. USA* 94:4113-4118). Many of the tau protein deposits also react with antibodies directed against ubiquitin. Immuno-electron microscopy demonstrates that tau is present as cytoplasmic filaments reminiscent of the paired helical filaments found in Alzheimer's disease, but these helical filaments have a different diameter and periodicity (Spillantini *et al.*, 1998, *Am. J. Pathol.* 153:1359-1363). Pathology varies between patients and may correlate with the particular *tau* mutation (Rizzu *et al.*, 1999, *Am. J. Hum. Genet.* 64:414-421).

Human tau protein has been expressed in transgenic mice in a number of studies. Gotz *et al.* (1995, *EMBO J.*, 14:1304-1313) used the human Thy-1 promoter to express, in transgenic mice, the longest isoform of wild-type human tau, which is 441 amino acids in length, containing the 58 amino acid amino-terminal insert and four microtubule binding domains. Brion *et al.* (1999, *Am. J. Pathol.* 154:255-270) used the murine 3-hydroxy-methyl-glutaryl CoA reductase (HMG-CR) promoter to express the shortest human tau isoform, which is 352 amino acids in length, contains no amino-terminal inserts, and has three microtubule binding domains. An increase in tau phosphorylation at sites associated with Alzheimer's disease pathogenesis is seen in both models, as shown by reactivity with the PHF1 and AT8 monoclonal antibodies that recognize phosphoepitopes on the tau protein. In the HMG-CR/*tau* mice, this increase in tau phosphorylation is accompanied by the formation of a conformation-dependent epitope associated with early Alzheimer's disease pathology (Brion *et al.*, *supra*). This epitope is detected with an antibody designated Alz50, raised against paired helical filaments purified from Alzheimer's disease brain. Neurofibrillary tangles were not present in either transgenic model.

Zahr *et al.* (1999, *Soc. Neurosci.* 25(A)447.1) have recently reported transgenic

expression in mice of mutant tau, containing the V337M mutation (*i.e.*, the valine at position 337 of the wild-type is replaced with methionine) in the context of a three microtubule binding domain isoform, under the regulatory control of the mouse prion promoter.

5 Additional transgenic models in which an increase in tau phosphorylation is seen include mice expressing mutant human amyloid precursor protein (APP) (Sturchler-Pierrat *et al.*, 1997, Proc. Natl. Acad. Sci. USA 94:13287-13292), glycogen synthase kinase-3beta (Brownlee *et al.*, 1997, Neuroreport 8:3251-3255), and the serine/threonine kinase Mos (James *et al.*, 1996, Neurobiol. Aging 17:235-241). Filamentous tau pathology is also
10 absent from these models.

Considerable effort has been put into making transgenic models of human neurodegenerative disease based on expression in transgenic mice of pathogenic human mutations. Choice of mutation, choice of transgene vector, expression in appropriate types of neurons, and transgenic expression of the mutant human protein at sufficient levels are
15 all critical to success. Strategies for creation of transgenic models have therefore varied widely. For example, the transgenic models for familial amyotrophic lateral sclerosis used the 12 Kb SOD1 gene for expression of human Cu,Zn superoxide dismutase containing the mutations G93A (Gurney *et al.*, 1994, Science 264:1772-1775) or G37R (Wong *et al.*, 1995, Neuron 14:1105-1116).

20 Transgenic models for amyloid β -peptide deposition in Alzheimer's disease have used either the platelet-derived growth factor promoter (Games *et al.*, 1995, Nature 373:523-527), insertion into the human prion gene (Hsiao, 1997, J. Neural. Transm. Suppl. 49:135-144), or the human Thy-1 promoter (Sturchler-Pierrat *et al.*, *supra*) to elicit transgenic expression of the amyloid precursor protein, containing either the "London" or
25 "Swedish" mutations associated with familial, early-onset Alzheimer's disease. These types of transgenic mice are becoming extraordinarily valuable to the pharmaceutical industry for their use as "gatekeeper" models for preclinical evaluation of promising drug candidates. An example of such a use was the testing of riluzole in the transgenic model of amyotrophic lateral sclerosis, where it proved to have a beneficial effect (Gurney *et al.*,
30 1996, Ann. Neurol. 39:147-157). The drug later became the first to be approved by the Food and Drug Administration for the treatment of the human disease. Similar use is

being made of transgenic mice that model various aspects of Alzheimer's disease pathology.

The current transgenic models of Alzheimer's disease, which are based on expression of mutant human amyloid precursor protein, are incomplete models of the disease. Although they show amyloid β -peptide deposition in plaque, no neurofibrillary tangles are noted, nor is there extensive neuronal cell loss or consequent brain atrophy. Thus, the art is in need of new and/or better models of Alzheimer's disease as well as other neurodegenerative diseases. The present invention is directed to addressing these, and other needs.

SUMMARY OF THE INVENTION

This invention provides transgenic mice expressing wild-type or mutant isoforms of human tau protein. The transgenic mice of the invention can be used for screening candidate drug compounds, which may be effective against neurodegenerative diseases and other diseases involving taupathologies. The transgenic mice of the invention can also be used in genetic crosses with other transgenic mice strains to model the progression of neurodegenerative disease.

In one aspect, the present invention is directed to a transgenic mouse expressing a human tau protein under the regulatory control of the mouse prion gene promoter. The human tau protein that is expressed in the transgenic mouse may be either wild-type or may contain a mutation. The mutation may be one previously identified in humans, or may be a novel mutation. The human tau protein expressed may be any of the six different isoforms.

The invention is also directed to the use of the transgenic mice of the invention as models of neurodegenerative disease.

In another aspect of the invention, the transgenic mice are used to discover drugs that modulate tau protein expression and/or activity.

These and other aspects of the invention are more fully described below.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows a schematic of a *tau* transgene construct used to prepare a

transgenic mouse of the present invention. See also Table 3, *infra*.

Figure 2 shows immunoblot analyses of human tau protein expression in the brains of wild-type and two kinds of mutant *tau* transgenic mice. Mouse brain samples are identified by mouse transponder ID number (5 digit number) or by ear tag number (337 and 341) (see Table 4, *infra*). Samples 21080, 21089, 21096, 22410, 22418, and 22435 serve as controls without human tau protein expression, and are from non-transgenic litter mates of the transgenic mice. Mice with transponder numbers are the progeny of the original founder mice mated with (C57Bl/6 x SJL)F1 partners; mice with three digit numbers are the founder mice.

DETAILED DESCRIPTION OF THE INVENTION

The invention provides transgenic mice, containing transgenes of human *tau*, and expressing human tau protein.

Transgenic animals contain within their genetic material some genetic information from another organism. For example, a transgenic mouse might have a human gene inserted into its genetic material. Such foreign genetic material is referred to as a transgene.

Transgenic mice can be generated in a number of manners known to those of skill in the art. Typically, a transgenic animal is generated according to the following basic steps: 1) engineer a nucleic acid construct with a promoter (regulatory sequences) and cDNA or gene sequences of interest; 2) linearize and inject the transgene construct into the pronuclei of embryos; 3) implant the embryos into the uterus of a pseudo-pregnant female; 4) screen the offspring that are born (the live pups in the case of mice) for the presence of the transgene; and 5) breed the transgene-positive pups (founders) to get transgenic "lines." Examples of these and related techniques are found in, *e.g.*, Hogan, B., Costantini, F. & Lacy, E., eds., *Manipulating the Mouse Embryo : A Laboratory Manual*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY (1986), which is incorporated herein by reference in its entirety.

Tau is predominantly expressed in neurons; the protein is principally localized in the axons. Some expression has also been noted in oligodendrocytes and astrocytes. Six different isoforms of tau protein are expressed in the adult human brain. Each isoform is

generated by differential splicing of an RNA transcript from the single *tau* gene, located on chromosome 17. The six isoforms range in size from 352 to 441 amino acid residues. The isoforms differ by having either three or four microtubule binding domains (imperfect repeats of 31 or 32 amino acid residues) in the carboxy-terminus, and by the presence or
 5 absence of 29- or 58-amino acid inserts, of unknown function, in the amino-terminus.

Tau protein microtubule binding domains, which contain the core microtubule binding domain motif proline-glycine-glycine-glycine (PGGG), are designated R1, R2, R3, and R4, and are encoded by exons 9, 10, 11, and 12, respectively. Exon 10, which encodes amino acid residues 275 through 305, is alternatively utilized (present in three of
 10 the six isoforms), such that the R2 microtubule binding domain is present only in tau isoforms containing four repeats. Thus, the isoforms range in size from 352 amino acid residues (with no amino-terminal inserts and three microtubule binding domains) to 441 amino acid residues (with a 58 amino acid amino-terminal insert and four microtubule binding domains). Table 1 summarizes the features of the six human tau isoforms, which
 15 are designated "A" through "F." The isoforms may also be respectively referred to as numbers "1" through "6," or by their total length in amino acid residues.

Table 1: The Six Isoforms of Human Tau Protein:

		<u>N-Terminal Insert</u>	<u>Microtubule Binding Domains</u>	<u>Length (amino</u>
20	<u>acids)</u>			
	A	none	3	352
	B	29 amino acids	3	381
	C	58 amino acids	3	410
	D	none	4	383
25	E	29 amino acids	4	412
	F	58 amino acids	4	441

In fetal brain, only the smallest isoform of tau ("A" in Table 1) is expressed. The amino acid residues in tau protein, regardless of isoform, are typically referred to by the
 30 position number they would have in the largest isoform ("F"), which has 441 residues. See Spillantini & Goedert, 1998, Trends Neurosci. 21:428-433.

The DFTP-17 associated mutations in *tau* provide an important entry point in the

study of neurofibrillary tangle formation as it relates to neuronal dysfunction and death. *Tau* mutations linked to FTDP-17 are either missense mutations (see Table 2), leading to substitution of one amino acid for another, or mutations in the 5' splice site of exon 10.

Most of the missense mutations associated with FTDP-17 occur in or near three of the four homologous microtubule binding domains within tau. Many of the missense mutations reduce the ability of tau to bind microtubules and to promote microtubule assembly in biochemical assays (Hong *et al.*, 1998, Science 282:1914-1917; Hasegawa *et al.*, 1998, FEBS Lett. 437:207-210). The arginine to tryptophan mutation at amino acid 406 (R406W), although linearly distant from a microtubule binding domain (residue 406 is encoded in exon 13, which does not contain a repeat), is believed to interfere with microtubule binding through a conformational change in the protein (Hong *et al.*, *supra*). The asparagine to lysine at 279 (N279K), proline to leucine at 301 (P301L), and serine to asparagine at 305 (S305N) mutations are seen only in four repeat-containing tau isoform proteins, as these residues are encoded by alternatively spliced exon 10. The valine to methionine at residue 337 (V337M) mutation is of particular interest, because it causes a tau pathology in the brain indistinguishable from that seen in Alzheimer's disease (Spillantini *et al.*, 1996, Acta Neuropathol. 92:42-48; Poorkaj *et al.*, 1998 Ann. Neurol. 43:815-825).

Table 2: *Tau* Missense Mutations Associated with FTDP-17

Codon*	Amino Acid Substitution	Location Within Tau Protein	Reference
272	Glycine → Valine (G272V)	R1 Microtubule Binding Domain	1
279	Asparagine → Lysine (N279K)	R1 Microtubule Binding Domain	2, 3, 7
301	Proline → Leucine (P301L)	R2 Microtubule Binding Domain	1, 2, 4, 7
305	Serine → Asparagine (S305N)	R2 Microtubule Binding Domain	3, 8
337	Valine → Methionine (V337M)	R3 Microtubule Binding Domain	5, 6, 7

406	Arginine → Tryptophan (N406W)	Carboxy-Terminal to R4 Microtubule Binding Domain	1, 7
-----	----------------------------------	--	------

* Numbering is based on the longest tau isoform, containing 441 amino acid residues.

1. Hutton *et al.*, 1998, Nature 393:702-705
2. Clark *et al.*, 1998, Proc. Natl. Acad. Sci. USA 95:13103-13107
- 5 3. Hasegawa *et al.*, 1999, FEBS Lett. 443:93-96
4. Dumanchin *et al.*, 1998, Hum. Mol. Genet. 7:1825-1829
5. Poorkaj *et al.*, *supra* (refers to codon based on numbering of a shorter tau isoform)
6. Spillantini *et al.*, 1998, Brain Pathol. 8:387-402
7. Hong *et al.*, *supra*
- 10 8. Iijima *et al.*, 1999, Neuroreport 10:497-501

Two studies have examined the effect of several tau mutants on microtubule structure after transfection of the *tau* genes into cells in culture (Dayanandan *et al.*, 1999, FEBS Lett. 446: 228-232; Arawaka *et al.*, 1999, Neuroreport 10:993-997). These studies show variable effects of the mutants on microtubule structure. Arawaka *et al.* (*supra*) showed that the presence of the V337M mutation results in fewer cell processes and reduced microtubule networks. Dayanandan *et al.* (*supra*) reported no differences in morphology with V337M, P301L or R406W, but did see differences in cell processes after treatment with cytochalasin B. In addition, Nacharaju *et al.* (1999, FEBS Lett. 447:195-199) have shown that mutants P301L, V337M and R406W cause an accelerated aggregation of tau into filaments *in vitro*.

A number of splice mutations have been identified in association with FTDP-17. These mutations are located in the 5' region of the intron immediately following exon 10 of the *tau* gene. The mutations are: G to A at nucleotide position +3 of the intron (Spillantini *et al.*, 1998, Proc. Natl. Acad. Sci. USA 95:7737-7741); A to G at +13 (Hutton *et al.*, 1998, Nature 393:702-705); C to T at +14 (Hutton *et al.*, *supra*, Clark *et al.*, *supra*, Hong *et al.*, *supra*); and C to T at +16 (Hong *et al.*, *supra*). These splice mutations all destabilize a potential stem-loop structure that may be involved in regulation of the alternative splicing of exon 10. This causes more frequent usage of the 5' splice site and

an increased proportion of *tau* transcripts that include exon 10 (Hutton *et al.*, *supra*). The increase in transcripts containing exon 10 presumably is mirrored by an increase in the proportion of tau protein containing four repeats of the microtubule binding domain. An increase in splicing of exon 10 is also seen with two of the tau missense mutations, asparagine to lysine at residue 279 (N279K) and serine to asparagine at residue 305 (S305N), neither of which alters microtubule binding or assembly (Hasegawa *et al.*, 1999, *supra*).

Based on what is understood about the mutations associated with FTDP-17, two types of mutations are sufficient to cause tau neuropathology: (1) mutations that alter the ability of tau to bind to microtubules and promote microtubule assembly; and (2) mutations that increase the proportion of tau protein containing four microtubule binding domains.

The present invention provides transgenic mice expressing human tau protein in either wild-type or mutant forms. In one embodiment of the invention, human tau protein is expressed in transgenic mice under the control of regulatory sequences from the mouse prion (PrP) gene promoter region. Preferably such expression is directed to the brain. More preferably, the regulatory sequences include the 5' flanking sequence of the mouse prion gene promoter, the first exon, the first intron, and the initial noncoding portion of the second exon of the mouse prion gene.

In a preferred embodiment, the human tau protein that is expressed in the transgenic mice is the wild-type 383 amino acid isoform of the protein (isoform D). This isoform contains four microtubule binding domains (*i.e.*, it utilizes exon 10), but has no amino-terminal insert; it is the shortest of the four repeat-containing isoforms.

In another embodiment of the invention, mutant human tau protein is expressed in the transgenic mice of the invention. Preferably, the mutation is the V337M mutation in the 383 amino acid isoform of the human tau protein. Those of skill in the art will recognize that other mutations can be used in the context of any of the six different isoforms of tau.

The tau protein expressed in the transgenic mice of the invention may contain a mutation known to be associated with FTDP-17, including, but not limited to, the G272V, N279K, P301L, S305N, V337M, and R406W missense mutations.

The invention also provides transgenic mice expressing mutant forms of human tau

protein, containing mutations other than those known to be associated with FTDP-17. Those of skill in the art will recognize that such mutations can be selected based on the locations of microtubule binding domains and other important features of the tau protein. Mutations to be incorporated into *tau* transgenes can be chosen based on predicted interference with tau function or analogy with disease-associated mutations. Such mutations can be tested for their effect on tau function by transfection studies on tissue culture cells. Those of skill in the art will recognize that any mutations in *tau* which modulate tau activity can be used in the transgenic mice of the invention.

Examples of amino acid residues in tau that can be mutated include residue 332, which is a proline in wild-type tau. Residue 332 is located in the core PGGG motif of the R3 microtubule binding domain. Mutation of residue 332 from proline to leucine (P332L), mimics the FTDP-17-associated alteration at residue 301 (P301L) in the core PGGG motif of the R2 microtubule binding domain. In a preferred embodiment of the invention, the mutant human tau expressed in the transgenic mice contains the P332L mutation in the 383 amino acid isoform of the protein.

A further embodiment of the invention includes transgenic mice expressing human tau protein containing a missense mutation at amino acid residue 364, where proline has been changed to leucine, resulting in alteration of the core PGGG motif of the R4 microtubule binding domain, also mimicking the FTDP-17-associated mutation, P301L.

In a preferred embodiment of the invention, the nucleotide sequence encoding the human tau protein is provided by a nucleic acid containing no intronic sequences. Other embodiments may include intronic sequences in the transgene to study the effect of mutations that occur in such intronic regions, for example, the mutation of G to A at position +3 in the intron immediately following exon 10. Thus, the invention also provides transgenic mice containing transgenes of human *tau* with intronic mutations.

In another embodiment, the transgenic mice of the invention are used as models of the progression of neurodegenerative disease or other diseases whose hallmark is neuronal cell death. Preferably, the transgenic mice of the invention will develop, with age, tau pathologies including tau hyperphosphorylation and filamentous aggregates of tau. Tau hyperphosphorylation may be revealed by immunostaining for tau phosphoepitopes such as those recognized by PHF1 or AT8 monoclonal antibodies. Filamentous tau neuropathology may be revealed by staining for ubiquitin immunoreactivity. More

preferably, the tau transgenic mice will display neuronal loss and cognitive dysfunction.

In another embodiment of the invention, transgenic mice expressing tau protein are used to screen for compounds that can be used to treat neurodegenerative disease.

It will be appreciated by those of skill in the art that the transgenic mice of the present invention are useful for analysis of neurodegenerative diseases involving tau pathologies. Such diseases involving tau pathology, collectively referred to as taupathies, include, but are not limited to, Alzheimer's disease, Pick's disease, FTDP-17, progressive supranuclear palsy (PSP), and corticobasal degeneration (CBD), Down's syndrome, Argyrophilic grain disease, amyotrophic lateral sclerosis/parkinsonism dementia complex of Guam, Non-Guamanian motor neurone disease with NFT, Niemann-Pick disease type C, subacute sclerosing panencephalitis, postencephalitic parkinsonism, dementia pugilistica, myotonic dystrophy, Gerstmann-Straeussler-Scheinker disease with tangles, prion protein amyloid antipathy, presenile dementia with tangles and calcifications, and Hallervorden-Spatz disease (Tolnay & Probst, 1999, Neuropathol. Appl. Neurobiol. 3:171-187).

Thus, in another embodiment, the invention is a model of neurodegenerative disease comprising any of the above described transgenic mice. In another embodiment, the transgenic mice of the invention are used in the screening of drug compounds to identify those capable of treating neurodegenerative disease.

In another embodiment, the transgenic mice of the invention are used in the screening of drug compounds to identify those capable of modulating the expression and/or activity of tau protein.

In another embodiment, the transgenic mice of the invention are used to screen for compounds that can be used to treat diseases that involve neuronal cell death.

Other embodiments of the invention will be readily understood by those of skill in the art.

The invention is further illustrated by way of the following examples, which are intended to elaborate several embodiments of the invention. These examples are not intended, nor are they to be construed, as limiting the scope of the invention. It will be clear that the invention may be practiced otherwise than as particularly described herein. Numerous modifications and variations of the present invention are possible in view of the teachings herein and, therefore, are within the scope of the invention.

EXAMPLES

Example 1: Preparation of Transgenic Vectors

Plasmid phgPrP, containing most of the mouse *PrP* gene 5' end and its 3' flanking
 5 sequences, was obtained from Marek Fischer, Department of Medicine, Division of
 Infectious Diseases and Hospital Epidemiology, University Hospital Zurich, Ramistrasse
 100, CH-809 Zurich, Switzerland. Plasmid T43, containing the cDNA encoding the wild-
 type 383 amino acid isoform of human tau [SEQ ID NO:1] in the vector pSG5
 (Stratagene, La Jolla CA; GenBank Accession AF013258), was obtained from Virginia
 10 Lee, Department of Pathology and Laboratory Medicine, University of Pennsylvania
 School of Medicine, 34th & Spruce Street, Philadelphia, PA 19104.

The sequence of the phgPrP insert was confirmed by the DNA Sequencing Core
 (Pharmacia & Upjohn). The 14,214 base pair (bp) phgPrP insert sequence was verified by
 comparison with the 38,418 bp GenBank sequence deposit for the *Mus musculus* short
 15 incubation prion protein *Prnpa* gene (Accession Number U29186). The phgPrP insert
 sequence extends the GenBank deposit by 5,047 bp of 5' flanking sequence [SEQ ID
 NO:2]. The entire phgPrP insert was cloned into pBluescript KS+ (Stratagene, La Jolla,
 CA). The resulting construct was opened with KpnI to remove the *PrP* coding sequence
 and 3' flanking region of the mouse *PrP* gene. Amplification by polymerase chain
 20 reaction (PCR) was used to resynthesize the 3' mouse *PrP* flanking sequence such that it
 contained 5' KpnI and XhoI sites for cloning. This created the vector 97-1, used for
 subsequent steps in assembly of the *PrP/tau* transgene.

To assemble the *PrP/tau* transgene, the 97-1 vector was opened with XhoI and
 Sall. This retained the mouse *PrP* promoter and sequences through the initial, noncoding
 25 portion of *PrP* exon 2. Sall was used to excise the human *tau*-383 insert fused to the
 polyadenylation site from the pSG5 vector. After dephosphorylation of the 97-1 vector,
 the two DNAs were ligated and transformed into *E. coli* strain DH5 α (GIBCO/BRL,
 Gaithersburg, MD). Bacterial colonies were screened for directional ligation of the *tau*
 insert in the sense orientation by PCR using the primers
 30 AGTAATTGAAAGAGCTCAGACGATG [SEQ ID NO:3] and
 TGTCACCCTCTTGGTCTTGGTGC [SEQ ID NO:4].

The V337M mutation was introduced into the *tau*-383 coding sequence by

mutagenesis of a 302 bp PstI-HindIII DNA fragment using PCR. The PCR reaction was carried out using: a primer positioned 5' of the PstI site having the sequence GTACTCCACCCAAGTCGCCGTC [SEQ ID NO:5], a short 3' reverse primer containing the HindIII site having the sequence GCAGCAGCATCGAAGCTTCTCAG [SEQ ID NO:6], and a longer reverse "patch" primer containing the V337M mutation having the sequence GCAGCAGCATCGAAGCTTCTCAGATTTTACTTCCATCTGGCCACCTCC [SEQ ID NO:7].

The P332L mutation was also introduced into the *tau*-383 coding sequence by mutagenesis using PCR. The PCR reaction was carried out using a forward primer having the sequence CCGCCAAGAGCCGCCTGCAG [SEQ ID NO:8], the SEQ ID NO:6 reverse primer, and a longer reverse "patch" primer containing the P332L mutation, having the sequence GCAGCAGCATCGAAGCTTCTCAGATTTTACTTCCACCTGGCCACCTCCTAGTTT ATGATGG [SEQ ID NO:9].

The PCR reaction was performed with an Advantage-HF™ high-fidelity PCR kit (Clontech, Palo Alto, CA), using the buffer supplied with the kit. The "patch" PCR primer was used at 1/100th the concentration of the short, flanking primers. Five cycles of (95°C for 45 seconds, 60°C for 2 minutes, 72°C for 3 minutes) followed by 20 cycles of (95°C for 45 seconds, 68°C for 3 minutes) were used for synthesis of the mutant fragment. To construct a mutant *tau*-383 insert, the PstI-HindIII fragment from wild-type *tau*-383 was exchanged for a mutant fragment. Proper introduction of mutations was confirmed by sequencing in the Pharmacia & Upjohn DNA sequencing core. The cDNA sequence for the V337M mutant of the 383 amino acid isoform is provided in SEQ ID NO:10. The cDNA sequence for the P332L mutant of the 383 amino acid isoform is provided in SEQ ID NO:11. The same procedure as described above for wild-type *tau* was used to assemble mutant inserts, *tau*-V337M and *tau*-P332L, into the *PrP* transgene vector. Table 3 presents the general features in the transgene constructs used to generate *tau* transgenic mice. The features correspond to those depicted in Figure 1. The sequence for the entire 9990 nucleotide sequence of the *PrP/tau* transgene vector comprising wild-type 383 *tau* is provided in SEQ ID NO:15.

Table 3. General Features in the *PrP/tau* Transgene Vectors Used for Microinjection

<u>Feature</u>	<u>Nucleotide Location</u>	
PrP promoter	1-6197	(Nucleotides 1-5047 are designated SEQ ID NO:2)
PrP exon 1	6198-6270	
PrP intron 1	6271-8457	
PrP exon 2	8458-8527	
<i>tau</i> -383 cDNA	8528-9727	
poly(A) region	9728-9990	

Example 2: Generation and Screening of Transgenic Mice

The 10 Kb plasmid inserts were prepared for microinjection by the University of Michigan Transgenic Facility following standard protocols. They were injected into F2 fertilized mouse eggs from matings of F1 hybrid C57Bl/6 x SJL mice. Progeny were genotyped by Taqman™ PCR using mouse genomic DNA prepared from tail biopsies using the Qiagen tissue kit according to manufacturer's instructions. Amplification was performed with the Taqman™ PCR Core Reagents Kit (PE Biosystems) using 50mM KCl, 10mM Tris-HCl, pH 8.3, 10mM EDTA, 60nM Passive Reference Dye 1, 5mM MgCl₂, 200μM each of dATP, dGTP and dCTP, 400μM dUTP, 200nM each primer, 100nM Taqman probe, 0.5 units uracil-N-glycosylase, and 1.25 units AmpliTaq Gold. Specific amplification of human *tau* DNA was performed using the primers G29-Fp GCATTGGAGACACCCCCAG [SEQ ID NO:12], G29-Rp GCTTTTACTGACCATGCGAGC [SEQ ID NO:13], and G29 Taqman™ probe with 5' and 3' colorimetric tags FAM-CTGGAAGACGAAGCTGCTGGTCACG-TAMRA [the nucleotide portion of which is designated SEQ ID NO:14] with the thermal cycling parameters, 50°C-2 minutes, 95°C-10 minutes, 40 cycles of [95°C-15 seconds, 60°C-1 minute], and 25°C hold. The starting amount of mouse genomic DNA template was 10ng. A standard curve was generated ranging 5 orders of magnitude from 10² to 10⁶ copies of linearized *PrP/tau* transgene plasmid DNA, which was diluted into mouse genomic DNA to give a final concentration of 10ng/μl. All reactions were performed in duplicate.

Multiple mice were generated from fertilized mouse eggs injected with wild-type

or mutant constructs. For each construct, several founder mice, carrying the *tau* transgene, were identified. Table 4 shows that the transgenic mice contained varying numbers of copies of the *PrP/tau* transgene, from which varying levels of transcript were expressed.

5 **Table 4: Summary of Analysis of Transgenic *Tau* Mice**

Transgene and founder mouse number	Mouse Id	Gene Copy #	<i>htau</i> RT-PCR levels copies/67ng total RNA positive RT sample	<i>htau</i> RT-PCR levels copies/67ng total RNA negative RT sample
TgN(<i>PrP/tau</i>) 355	21088	4.6	16,599	81
TgN(<i>PrP/tau</i>) 327	21097	2.6	5,402	13
TgN(<i>PrP/tau</i>) 326	21083	13.7	595	161
	21214	10.6	754	26
TgN(<i>PrP/tau</i> -V337M) 249	22419	2.4	33,671	6
	22420	2.6	26,912	20
TgN(<i>PrP/tau</i> -V337M) 250	22405	0.3	19,138	0
	22408	0.2	21,814	0
TgN(<i>PrP/tau</i> -V337M) 257	22436	0.9	10,755	2
	22437	0.2	8,842	1
TgN(<i>PrP/tau</i> -P332L) 337	337	16.8	ND ¹	ND
TgN(<i>PrP/tau</i> -P332L) 341	341	1.4	ND	ND

¹Not determined.

10 Table 4 summarizes the analysis of the transgenic mice. The first column lists the transgene designation (TgN) and founder mouse number. The founder mice are indicated by the numbers 355, 327, 326, 249, 250, 257, 337 or 341, after the transgene name TgN(*PrP/tau*), TgN(*PrP/tau*-V337M) or TgN(*PrP/tau*-P332L). The second column lists the mouse identification (Id) number (either a 5-digit transponder number or a 3-digit founder number, as used in Figure 2). Mice with a 5-digit number are the progeny of the founder mouse indicated in the first column by the three digit number after the transgene

name. The third column lists the haploid copy number of the human *tau* (*htau*) transgene, based on DNA levels derived after PCR analysis. This number was measured either in the founder or founder progeny, as indicated in the second column. The fourth column lists the human *tau* (*htau*) transgene mRNA expression, as determined by reverse transcriptase PCR (RT-PCR). The fifth column lists the background amplification of the RNA samples, without conversion to cDNA by RT.

Example 3: Human Tau Expression

Mouse brain samples were harvested after intracardiac perfusion with cold phosphate-buffered saline, for analysis of human *tau* mRNA and tau protein expression. For cardiac perfusion, mice were deeply anesthetized, then perfused transcardially until the effluent was cleared of blood. Tissue samples were frozen immediately on dry ice, then stored at -70°C until analysis.

Total RNA was prepared from homogenized cortical brain tissue using the RNeasy mini kit (Qiagen). The RNeasy protocol was modified by the addition of the RNase-free DNase set protocol (Qiagen) to treat the RNA, while bound onto the column, with RNase-free DNase I at room temperature for 15 minutes to 1 hour, to remove any residual genomic DNA contamination. Two micrograms of total RNA was converted to cDNA in a 30µl reaction containing, 50mM Tris-HCl, pH 8.3, 75mM KCl, 3mM MgCl₂, 500µM each dNTP, 0.5µg random hexamers, 30 units RNase inhibitor, and 400 units Superscript II reverse transcriptase (RT), which was incubated at 42 °C for 1 hour. The Superscript II RT enzyme was then heat inactivated by incubation at 95°C for 5 minutes. To assess the extent of genomic DNA contamination in the total RNA samples, control reactions, without RT, were also performed for each sample.

For determination of human *tau* mRNA expression, PCR amplification on the ABI PRISM 7700 instrument was performed using 2X Taqman™ universal PCR master mix (PE Biosystems) containing 200nM G29Fp and 200nM G29Rp primers and 100nM G29 Taqman probe. The final PCR reaction volume was 25µl, and contained 5µl of a 1:5 dilution of the cDNA reaction described above. The thermal cycling parameters were as follows, 50°C-2 minutes, 95°C-10 minutes, 40 cycles of [95°C-15 seconds, 60°C-1 minute], and 25°C hold. A relative standard curve was generated ranging 5 orders of magnitude

from 10^2 to 10^6 copies of linearized *PrP/tau* transgene plasmid DNA, which was diluted into mouse genomic DNA to a final concentration of 10ng/ μ l. All samples were run in either duplicate or triplicate.

For determination of human tau protein expression by immunoblot analysis, brain
5 tissue was minced in 600 μ l lysate buffer (0.1M MES, pH 6.8, 0.75M NaCl, 0.5mM EGTA, 2mM DTT, 2mM PMSF, 10 μ g/ml aprotinin, 10 μ g/ml leupeptin, 10 μ g/ml soy bean trypsin inhibitor, 5 μ g/ml pepstatin) on ice. Brain fragments were ground with 10 strokes of a pestle in a pre-chilled TenBroeck tissue homogenizer. Lysate was centrifuged at 40,000g for 15 minutes at 4°C. Protein concentration of supernatants was determined by
10 Bio Rad protein assay, and lysates were stored at -80°C until used.

50 μ g samples of the brain lysates were diluted into 4X gel loading buffer (250mM Tris-HCl, pH 6.8, 8% SDS, 20mM EDTA, 40% glycerol, 0.02% bromophenol blue, 20% β -mercaptoethanol), heated at 100°C for 5 minutes, and electrophoretically separated on a
15 7.5% polyacrylamide gel in running buffer (25mM Tris base, 192mM glycine, 0.01% SDS). The proteins were transferred to Nitrobind nitrocellulose membrane (Micron Separations Inc, Westborough, MA) in transfer buffer (25mM Tris base, 192mM glycine, 20% methanol). The membranes were blocked for 1 hour in PBS containing 5% nonfat dry milk. Blots were probed overnight at 4°C with either the T14 or HT7 anti-human tau antibody (Zymed Laboratories, Inc., So. San Francisco, CA), diluted to 2 μ g/ml in a
20 solution of 10mM Tris-HCl, pH 7.5, 500mM NaCl, and 0.1% Tween 20 (TNTween), plus 1% BSA and 3% dry milk. After washing in TNTween, the blots were incubated for 1 hour at room temperature with goat anti-mouse, horseradish peroxidase-conjugated secondary antibody (Kirkegaard and Perry Laboratories, Inc., Gaithersburg, MD) at 1:5000 dilution. The immunoblots were washed again in TNTween, treated with ECL
25 reagents for 1 minute, and exposed to X-ray film.

Progeny of the transgenic founders (mated to F1 hybrid (C57Bl/6 x SJL) partners) or the founders were analyzed for *PrP/tau* transgene expression in the brain. Both human *tau* mRNA (Table 4) and protein (Figure 2) were detected. Efficient, relatively high levels of human tau protein expression were obtained in transgenic mice using a mouse *PrP* gene
30 vector coupled to a heterologous human *tau*-383 cDNA. Roughly equivalent levels of protein expression were obtained from wild-type and mutant *PrP/tau* transgenes,

suggesting that expression of mutant tau protein is relatively well tolerated in the mouse. The levels of expression achieved appear to be higher than those reported by Gotz *et al.*, *supra* or Brion *et al.*, *supra*, although different antibodies and methods of analysis were used to establish levels of protein expression. Ranking by protein expression roughly
5 correlates with the PRISM measurement of mRNA expression.

Example 4: Microtubule Organization in GFP-*tau* Transfected Cells

To investigate the effect of mutations in the microtubule binding domains on tau protein interaction with microtubules, human embryonic kidney HEK293 cells were
10 transiently transfected with cDNA encoding the 383 human tau isoform fused to green fluorescent protein (GFP). Wild-type and the P332L, V337M, and P301L *tau* mutants were analyzed. Fluorescence and biochemical analyses showed that GFP-labeled mutant tau proteins P332L, V337M, and P301L have an altered intracellular distribution, as compared to GFP-wild type tau protein.

15 Studies using an anti-tubulin antibody revealed that GFP-wild-type tau protein co-localizes with microtubules. Some tau protein is also present in the cytoplasm, not associated with microtubules. In cells expressing high levels of wild-type tau, a reorganization of the microtubule cytoskeleton was observed. Bundles of microtubules could be seen, often in the form of a spiral or a ring around the circumference of the cell.
20 This is a phenomenon previously noted in non-neuronal cells expressing tau. Cells expressing low levels of GFP-wild type tau protein, contained a fine network of microtubules, often emerging from microtubule organizing center.

Cells expressing GFP-labeled mutant tau proteins P332L, V337M, and P301L exhibited higher levels of fluorescence dispersed in the cytoplasm than did cells
25 expressing GFP-wild type tau protein. The microtubule bundling and the microtubule network, seen in GFP-wild type tau expressing cells, was also reduced in cells expressing the mutant tau proteins.

Immunoblotting analyses of cytosolic and cytoskeletal cellular fractions confirmed the fluorescence studies, showing that cells expressing the GFP-mutant tau proteins
30 contained more tau protein in the cytosol than did cells expressing GFP-wild type tau. The difference was most dramatic for P332L, but the studies have confirmed that these

microtubule binding domain mutations interfere with tau protein interaction with microtubules.

Example 5: Cellular Analysis of Tau Transgenic Mice

5 General pharmacokinetic studies are used to analyze human tau expression in the transgenic mice. Pharmacokinetic studies include determination of the brain distribution of tau protein expression, and kinase activity assays. The state of phosphorylation of the tau protein is detected using monoclonal antibodies to specific phosphoepitopes of tau, including PHF1 and AT8, and various polyclonal antibodies. Filamentous tau
10 neuropathology is revealed by staining for ubiquitin immunoreactivity.

Example 6: Analysis of Neurodegenerative Disease in Transgenic Mice

Neurodegenerative disease is studied in *tau* transgenic mice by use of a variety of behavioral studies, including the Morris water maze test, mobility and gait tests, and
15 behavioral observation (see, e.g., Moechars et al., 1999, Biol. Chem. 274:6483-6492).

Example 7: Drug Screening in Transgenic Mice

The transgenic mice are used for the screening of compounds that affect tau hyperphosphorylation, filament formation, or neurodegenerative disease criteria.
20 Compounds that inhibit kinase activity, protease activity, phosphatase activity, oxidative stress reactions, apoptosis, or protein aggregation are tested for their ability to modulate 1) tau phosphorylation, 2) tau aggregate formation, 3) formation of cellular inclusions of paired helical filaments of tau, and 4) formation of neurofibrillary tangles in the transgenic mice. Those compounds exhibiting positive effects in such tests may be useful in the
25 treatment of neurodegenerative diseases.

All references cited herein are incorporated herein in their entirety by reference.

We claim:


1. A transgenic mouse, comprising a transgene, said transgene comprising a polynucleotide encoding a human tau protein operably linked to at least a portion of a regulatory region of a mouse prion gene, wherein said transgenic mouse expresses human tau protein.
5
2. The transgenic mouse of claim 1, wherein said human tau protein is selected from the group consisting of isoforms 352, 381, 410, 383, 412, and 441.
10
3. The transgenic mouse of claim 2, wherein said human tau protein is isoform 383.
4. The transgenic mouse of claim 1, wherein said human tau protein is selected from the group of mutants consisting of G272V, N279K, P301L, S305N, V337M, and R406W.
15
5. The transgenic mouse of claim 4, wherein said human tau protein is a V337M mutant of isoform 383.
20
6. The transgenic mouse of claim 2, wherein said human tau protein is further selected from the group of mutants consisting of G272V, V337M, and R406W.
7. The transgenic mouse of claim 3, wherein said human tau protein is further selected from the group of mutants consisting of N279K, P301L, and S305N.
25
8. The transgenic mouse of claim 2, wherein said human tau protein is mutant P332L.
30
9. The transgenic mouse of claim 1, wherein said human tau protein is expressed in the brain.


- 5
10. The transgenic mouse of claim 1, wherein said regulatory region comprises said prion gene promoter and 5' flanking sequence, the first *PrP* exon, the first *PrP* intron, and the initial, noncoding portion of the second *PrP* exon.
11. A model of neurodegenerative disease comprising a transgenic mouse as in any of claims 1-10.
- 10
12. The model of neurodegenerative disease of claim 11 for use in the screening of drugs to treat said disease.
13. The model of neurodegenerative disease of claim 11 for use in genetic crosses to generate models of Alzheimer's disease.
- 15
14. A method of screening for a drug that modulates hyperphosphorylation of tau comprising the steps of
- a) administering the drug to a transgenic mouse, the transgenic mouse comprising a transgene, said transgene comprising a polynucleotide encoding a human tau protein operably linked to at least a portion of a regulatory region of a mouse prion gene, wherein the transgenic mouse expresses hyperphosphorylated human tau protein; and
- 20
- b) comparing the state of phosphorylation of tau in a second transgenic mouse to which the drug was not administered to the state of phosphorylation of tau in the transgenic mouse to which the drug has been administered, wherein a difference in phosphorylation indicates the drug modulates hyperphosphorylation of tau.
- 25
15. A method of screening for a drug for treatment of a neurodegenerative disease comprising the step of administering the
- 30

drug to a transgenic mouse which exhibits neurodegenerative disease characteristics, the transgenic mouse comprising a transgene, said transgene comprising a polynucleotide encoding a human tau protein operably linked to at least a portion of a regulatory region of a mouse prion gene, wherein the transgenic mouse expresses human tau protein, and wherein it is determined whether the drug at least partially abates at least one of the characteristics of the disease.

- 10 16. A method of screening for a drug that blocks hyperphosphorylation of tau comprising the step of administering the drug to a transgenic mouse, the transgenic mouse comprising a transgene, said transgene comprising a polynucleotide encoding a human tau protein operably linked to at least a portion of a regulatory region of a mouse prion gene, wherein the transgenic mouse expresses hyperphosphorylated human tau protein, and wherein it is determined whether the drug at least partially blocks hyperphosphorylation of tau in the transgenic mouse.
- 15
- 20 17. A method of screening for a drug that blocks formation of filamentous aggregates of tau comprising the step of administering the drug to a transgenic mouse, the transgenic mouse comprising a transgene, said transgene comprising a polynucleotide encoding a human tau protein operably linked to at least a portion of a regulatory region of a mouse prion gene, wherein the transgenic mouse expresses human tau protein forming filamentous aggregates, and wherein it is determined whether the drug at least partially blocks formation of filamentous aggregates of tau in the transgenic mouse.
- 25
- 30

Figure 1



2/2
Figure 2

SEQUENCE LISTING

<110> Pharmacia & Upjohn Company

<120> Transgenic Mouse Model Of Human Neurodegenerative Disease

<130> PHRM0301

<150> 60/177,319

<151> 2000-01-21

<160> 15

<170> PatentIn version 3.0

<210> 1

<211> 1152

<212> DNA

<213> Homo sapiens

<400> 1

atggctgagc cccgccagga gttcgaagtg atggaagatc acgctgggac gtacgggttg	60
ggggacagga aagatcaggg gggctacacc atgcaccaag accaagaggg tgacacggac	120
gctggcctga aagctgaaga agcaggcatt ggagacaccc ccagcctgga agacgaagct	180
gctggtcacg tgacccaagc tcgcatggtc agtaaaagca aagacgggac tggaaagcga	240
gacaaaaaag ccaagggggc tgatggtaaa acgaagatcg ccacaccgcg gggagcagcc	300
cctccaggcc agaagggcca ggccaacgcc accaggattc cagcaaaaac cccgcccgc	360
ccaaagacac caccagctc tgggtgaacct ccaaaatcag gggatcgcag cggctacagc	420
agccccggct ccccaggcac tcccggcagc cgctcccga ccccgccct tccaacccca	480
cccaccggg agccaagaa ggtggcagtg gtccgtactc caccgaagtc gccgtcttcc	540
gccaaagacc gcctgcagac agccccctg cccatgccag acctgaagaa tgtcaagtcc	600

2

aagatcggct ccactgagaa cctgaagcac cagccgggag gcggaaggt gcagataatt 660
 aataagaagc tggatcttag caacgtccag tccaagtgtg gctcaaagga taatatcaaa 720
 cacgtcccgg gaggcggcag tgtgcaaata gtctacaaac cagttgacct gagcaaggtg 780
 acctccaagt gtggctcatt agscaacatc catcataaac caggagggtgg ccagggtggaa 840
 gtaaaatctg agaagcttga cttcaaggac agagtccagt cgaagattgg gtccctggac 900
 aatatcacc cgtccctgg cggaggaaat aaaaagattg aaaccacaa gctgaccttc 960
 cgcgagaacg ccaaagccaa gacagaccac ggggcggaga tegtgtacaa gtccgacgtg 1020
 gtgtctgggg acacgtctcc acggcatctc agcaatgtct cctccaccgg cagcatcgac 1080
 atggtagact cgtcccagct cgtcacgcta gctgacgagg tgtctgcctc cctggccaag 1140
 cagggtttgt ga 1152

<210> 2

<211> 9990

<212> DNA

<213> Mus musculus

<400> 2

ggcggccgag acggatccaa aggcagcaaa aaggcagaga ggggtgatact gggcctggct 60
 taagcatttg aaacttcaaa gctcaccccc aattacacac ttcttccaac aagtcacac 120
 ctctaatta gtccactct ctgtgggcct acggagagta ttttcttct aactaccaca 180
 gttgctgagg aatttaatta aaactacaac cttatcccaa cctagatctt tcagcctttc 240
 tgtactacca gagaggggtc atacagcatt gttgtgactc ccattataac ttaaaggga 300
 gctcacacaa agtccagagc cctccatacc ctgcaaatga agaagtacgt tctcaaatcc 360
 cttggagcag ggtccactt tggcggcaca aactttaatt tctagacgga acggcatctc 420
 tacagaaaga aaagccatgg tatctgcatg ataagtctga aaaggacctg ggcaaactctg 480
 cagctgacaa ttccagccat tgctgccact gcgagaaaac cctgctgatg gcagcattgt 540
 cagcatcatc tctccagga acaccggcca tcgagccacg aggacaattg ctgctgctgg 600
 agtcaattca tctgccagcc acatcatact ctgggaccgt cactaaccag atccaagcag 660
 ccttgaggaa gcatgtcttc tgggtgtgac tgatcccaag ggctgacaac aaggctctca 720
 cagaggcatc ttatgtcaac ctatctacca tgcacggtat aagacacatt ctctctgtg 780
 ctgtgtggac actgccatca cagcaacag aaaggaaact cactcactgt gtctgatgtg 840

gtggtgcttg ttaggggagt tctgggcatg tatggcacca tcgcccata ggactcctgt 900
ggggtcatgc ccactctact cctctagaga ccatgaagag atggagaggg aagagcaagc 960
acagatgaca ggctagaact aaagaggagt gtcaggtag cggacctgaa ctcacggctg 1020
ctcagcctga agtggtaggg ccactctcat ctggtatctg gctgaaggt gcgtggatac 1080
cctctgtgcc cgtccagaag ttctctactg aagacagaaa tgcctgtcca gtcattggaag 1140
aatgattggc agtccctact tctcagacca ctgaatgggt cagaacaact actgggtgac 1200
cctaaggtag tcttcagcag atatgtgtga aaaatggaaa gaagatgggt agaaataaac 1260
ggtttttagag gaaaaaaact ctcacaaaga tattataaaa agaaaagagc ttattatttg 1320
agcaagcatt caaccagaat gcacaccaca ggcagtctgc taaggagtg tgcagacagg 1380
aggagtgtcg ccccttatgt gagccagtag ataaggatgc tgtgctgtt tttagtaact 1440
ggctctcagc ttgacagcac cttttatcac atggtttaac cttaaattcat ctggcgaatg 1500
aggctgtcac gtacttctg attagcttta tctgaaatga gacaagcttc acatgttcac 1560
ggcaggaggt aatcctgctg cttagagaac agggccatc caagccaggc tcctctctcc 1620
accaacacgg gtggttgaag agctatctct cctggtgtg tgtgtttcag agatggctcc 1680
caggtttttg gtttggttg aattgggttt tggttttctt actctagccc agactagctt 1740
ggaattctct ggaaagctgc aacggggagc tcaggttcag tgagagatcc tgtctcaaaa 1800
agcagggtga gaagtgttg aggaagacac cccagtgtta acctctgacc tccatatgtg 1860
catgcatgga cagcatgga tacacataca cacacacaca cacacacaca cacacacaca 1920
cacacacaca cacacacaaa accagaaaga atgaacgccc cctctccagc ttgtttacag 1980
tagatacaga gcactcgtaa aacatggggg gtaaaactgaa tgctgagagt aacttagatg 2040
agtaattaag gaaggaagag gaaagaaacc aggaaccga gagcaagtga ctggaagatc 2100
gttaggcaat cccacacccc tgctcgttga agttggaatg cttctctctt ctgctcttg 2160
aagttcttta gaagtgttag gatctcaca ttagtctgtg gtggtttcaa tatgcttcac 2220
ccgtggttag tggcactatt aggaacgtg tcctgttga aggaagggtg tcactgcata 2280
ggcgggcttt gaggtgtctt ccagtgtca agctctccc agtgcaagag aggcagacac 2340
ctgttgctg cagaagacag tctctgctg cctttgaatc aagatgtaga actcaagccc 2400
catgtctgcc tgaacctctg aaactgtaag ccagcccaa ttaaatgtt tctttcaca 2460
gagttgcctt ggtcatggtg tctgttcaca gcaataaaac cctaactaag acagtcttaa 2520
atcaatgaaa agacctttaa ttattcattg aacaaacacc atttctctgt atcaagttgg 2580
cagtgactag taagcaacta tagttctgca ccagggacct ttttgagaa atataccgat 2640

ccaagcatgt tggcatctag attccaaagc caagacacct gccacacctt tccatgcctt	2700
gggttcctgg cagggcatct ggcttcgggg atgtgtatc caggcaccga ctggaatgca	2760
tggaaacaat taaaatagca tcatagaaga cattgcaatc ctaggagaga actataccaa	2820
aactcagaac tatacctggt taagtgtaga aaagacgaaa ggaataaaaac caggaatatt	2880
ttaaaatatt ttatttgagc tcatgtgcat gggatatttg cctgaaagta tgtctgtgta	2940
ccacatgcat ggctggctcc tgcagaggcc aaaagagagc atcagatctc cttagagctgg	3000
agtttcagaa gtttgtgagc taccacatgg gtgctggaaa caaaaccag gacatctgga	3060
agagcagcca gtgttcttaa ctactgagcc atcaccagg tcccaccatg aatgtttttc	3120
tttattcttc tctatatttt ctaatgtttt tattggaaat atacaacttt tgccacacat	3180
aacaaatgac caaagaaatg aggtgagagg ggcagctgtt caaatgctgc ctgggaaggc	3240
ttggccagcc ctggcttggc tgccctggc tcagctggcc ctgacttggc tgtccgggtg	3300
ccagctgtca tctactgctt cataataagc tgcactttgg gctgaagggg tggctcagcc	3360
tttaaaggct aggtcataa ccaaagtaag ttgcatttta ttgactag gttgaagggg	3420
gatctgaaac ttgctgtcaa tgttataaaa cattttatct tcaaatttgg tataggggtc	3480
atagaccaa ggttctataa accccagaac agcaccactc cctagaaata agcaccata	3540
caagagccta tgggacactt tatagccaaa caaaaagcta tgtttgaaac ttcctttaca	3600
agggcctgag tcccatctat aagggaagga gcccacttc gtaataacac cccactggtg	3660
acatttgaag gggacacatt caaactgtaa caccatctta tatcatttgc acattagggc	3720
caaactgtgc cagttgtca tttctaagaa gacagaagtt gtcaagcctg tgctttgagc	3780
cacaagtgtg acaacctact ttcaggcaag tcgctacttc cctaagactc taccccaata	3840
ggcctggggc ctggaatgtg ttaacacag atgcaggctt ctgccttagt gcaggcttga	3900
gttctcatgt cctctctctt ttagctttcc gtctcaagge gcctctcctt agcagaaaaa	3960
atcagaggca taaagcatac atcaggggga agccagagtt ttcagaggga gttttgtgat	4020
ggccttttca gagcattctt gtcaagacta gtttgctcg ttctctttat taaatgaaag	4080
aaaaataatg cagtgttgca aattagcttt ggtaatggct ccaaccattg tcagggttcac	4140
agtctcatc cgccattcaa aacaacaaac ccaccacact ctctatgcag tgccgtaact	4200
cagaacagcc accaaacagc agaaagagge tcccgcactc ctctcagcct tgccataaac	4260
tcgccggcca catgcttatt ttaaattatt taaattatgt cgtttctccc aacaatgacc	4320
tcccaagtgc ttggttgaca ggcttatacc attagccga ggcttgcata gcaacgataa	4380
ccaggtaggc tattattata accaggtagc tgccgagcta ctggtcgggc cctttttgtc	4440

5

tctagaaacc tctcaacccc cacccaaaaa agctttttatt gccacttcct agtgggtaga 4500
 gagcagtcag ccaatagata tttgattctt tgaggaaaaa gctgagtttt gatgtctttt 4560
 aatcaagcct ttcagagtcc ctctgtgggg gaggccaggt ggaagcgggg tgggaagctg 4620
 gtcccttacc taagctaata tagacacct cccactctc ccctgccctc ttgacagatg 4680
 cagtcatcct gatcacaatg agtattctct gaggcaggaa ggcaaggctc tgggaagatgg 4740
 tcaatgcctt cattaagaac ccagagtaaa ggtcaagcag acaccagcac cgcgaaatc 4800
 taatttcact gtaattgaat catctcagcc aaaggctgta ttttcagcc ctctcgtggc 4860
 ctcttcccca acaactgtca acaactgtgt gagcctacct atgtatgcgc gctcacacac 4920
 acacacacac acacacacac acacacacac aggggtggggg gacacaatga ttacacaaga 4980
 gtacttaata aacaactata attctcctgg ctcggaatgt tccctaccac cctctcctcc 5040
 tggatccgga tccataact ggatacaaat atttaataca aaccaatct tgtgtctgtt 5100
 aatgatcttc agtgtctcgc cctcagcaag aggacaggat attatgtttt cctgtgatt 5160
 tatgacctct tctgtctcag tatcggcagc aatttattta catggctttg gactgtgtta 5220
 tatgtgtagt atggacatga ggggtgcatgt caacctatgt gtggaggcca gaggtcaatg 5280
 tcatgtcttc cccaatcact gccagtggt ccctggattc caaactcagg tcttcacgct 5340
 tgggaactga gccagtgcc cagctcctaa ccctcccctg ttttaaaaag gtctcattat 5400
 gttgccagg tcagccttga acttgagagt ctctgactg caggcttca cctgtccaag 5460
 tcagcaggca tctgaacaa gaacatcatt tcccttaagc tgtttcaggc tgtgtttggt 5520
 gggagctgtt aaatgcagtg cattttttcc tttggacaca ataaaagaaa aaagtgatta 5580
 aatgagttgg gtgtggtggt gcgagtctac aatcctagaa ctgaggagat tgaggagaaa 5640
 gcattgctct gattttgagg tcagcttaaa ttacttagta ggaacaccag gccaaattgg 5700
 gctatgggat tgtctccaaa gataaagaaa aaaggaagg agagaaaaga aaaagaaagg 5760
 aaagaagggg aaaagaagga atcagcagag aataaataag tcaacatgca atggccaata 5820
 tactttctag gcctctaatt cttttatagt ttgtgggaaa atgtcgaaaa tcttcgttac 5880
 caatttcttg ttaccaaagt tcaacgatgg ctctctcgt cgttaggta acctttcatt 5940
 ttctcaacta cccattatgt aacgggagca ttgggtactg gatcagtcct ccattaaaga 6000
 tgatttttat agttgctgag cgtcgtcagg gactgctgac actgggggag gtttaaacag 6060
 atacaagcat ttaagccagt ccggagcggg gactcatccc ccccccccc ccccccccc 6120
 cgagagacgc ggcgcggcca ttggtgagca tcacgccccg cccctcgccc cgcctagttc 6180
 ccgcctgcc cgcctcttc cactcccggc tccccgcgt tctcgatca gcagaccgat 6240

tctgggcgct gcgtcgcatc ggtggcaggt aagcgggctg ctgaagccag gccttggcga 6300
 gcactcagcc ttcgcgtcgc aagcgggct cactgcgcct ctgggggcct tgaggccacg 6360
 gggactagga ctgggactgg gactggggct gagtctggct gggagggtgac tgtacacccc 6420
 ctgctgcgcg actcctggag gaaccgaatc ccagggcagc caggccggga gccagccttt 6480
 ccttcccagag ccagattcac agctcagcat cgctggggat ggggggtggca tcttttgact 6540
 gtccttggct gttttcttct ctctttagtag tagctacagc gaacataatt ttacctcgtt 6600
 attccaccac agtcattact cccttcgaca gtttcattct caacgtcgcc gtgcgccttc 6660
 actgcctgt ctaggcgttt tcatgattgt ctattttctt gtactttgaa taccgtgggt 6720
 taatagcagt tgcgggtgcg cagaattctc catttcctta agagaaactc ctgggagaat 6780
 gggactaaag acgtgcaaatt ttaattatat cgcaaacagg aatcaaaatt ttgcattaaa 6840
 atgccaaaca tcttgaaaaa ttaactattc aatgaagaaa aggaactact ttacctacac 6900
 acacatccga gagcttcgag gaggcgaagg aatagaaaag ctaagggatg atttgggttg 6960
 tatttgaatc tgacacaagc tttccatatt atttatagca gggactaaac gatgagtcac 7020
 tttctgaata agatgcaaatt taaagcaagt ttgtttgttg tctttacatc tattaatatg 7080
 acagagacaa tggcaacagc aaccctaacc tagaggttgc ctgaaagtgt caggtttggg 7140
 aacaagtggc cctgcttaag ggctagaaaag attgctttac aaccaacaat catgacttga 7200
 cattgcctgg ggttcctttt gtctattcct tttttaaag actagtgtt attttatgtg 7260
 catgagtgtt ttgcatccac attgcctgt atacacacct ggttctgtgg aggtcaggag 7320
 aggtgtgtgg atgcctggc actagagcct tggatggtta tgtgagcccc tgccacaggg 7380
 gagctcagaa ccaatccag gtccctctgga agagcaacca gagctcttaa aacttctaag 7440
 tatccctcca tcccctttcc atcatatttg gaaaggagaa aactgctacc catgcctggc 7500
 atttatttca gagattaact gtctgtgtaa aacttgacat tgaaagtgca ctattctgtt 7560
 tcccattcat acttagttga gactactgta agtcagttag ggctttttt gtttggttcc 7620
 ttggttagtt tggagtgtgt ttgtgagctc attaacaggc tttcaatatg tagctggaat 7680
 ttgctgtgta gaccagacag gcctcaaatt tgtggcaatc ctccctgcat cttcccagaa 7740
 tgccctggta caggcataaa ccaccgtgcc cagcagtaaa acaatctggg gaggtattat 7800
 tagtcgtgtg ctgtgacca gaaacccac tcctggcaat ttactgggaa ggaacaaaca 7860
 aagggttagg ggagccatat ggcctgcagt tagagaaaat tagatccaac tgaaaaatca 7920
 acctaaagggt gtaaaagcca agcagttaag aaactgacaa gctcatgatg gaagccgagg 7980
 ccatcgtgaa cactcttcat tttaggcccc acgtatcact ggggacaact gagagtcaaa 8040

gtacaggtaa ggagaccaag gcttttcagg actcaggctg tctcagtga aagcccagaa 8100
 gagcagtaat tgaaagagct cagacgatgt gtctgatctc ccttggttgt ttgttgcgtg 8160
 attatttcca ctaacttatt tgggaggaaa aaaaacagtt cacaggcttc ttttcttgaa 8220
 atactgggga ttgctgggat cgaaccagg gataggtttt tagtttctaa aataacatag 8280
 atcatgccct gtttgctttt tggaaatatgt ttgcgctgcc cttattttca tgttcaaata 8340
 ctgctccatt ttgcgtgact ctttagtatt ggtttgatga tttgcatatt agattagatt 8400
 gtatttcagt tctcagactt atttatcaat tctagttttc tctttttgtt gttttaaagg 8460
 actcctgagt atatttcaga actgaaccat ttcaaccgag ctgaagcatt ctgccttcct 8520
 agtggtagct cgactatcag gtgaactttg aaccaggatg gctgagcccc gccaggagtt 8580
 cgaagtgatg gaagatcacg ctgggacgta cgggttgggg gacaggaaaag atcagggggg 8640
 ctacaccatg caccaagacc aagagggtga cacggacgct ggcctgaaaag ctgaagaagc 8700
 aggcatgga gacaccccca gcctggaaga cgaagctgct ggtcacgtga cccaagctcg 8760
 catggtcagt aaaagcaaag acgggactgg aagcgatgac aaaaaagcca agggggctga 8820
 tggtaaaacg aagatcgcca caccgcgggg agcagcccc ctaggccaga agggccaggc 8880
 caacgccacc aggatccag caaaaacccc gcccgctcca aagacaccac ccagctctgg 8940
 tgaacctcca aaatcagggg atcgacgag ctacagcagc cccggctccc caggcactcc 9000
 cggcagccgc tcccgacccc cgtcccttcc aacccccccc acccgggagc ccaagaaggt 9060
 ggcagtggtc cgtactccac ccaagtcgcc gtcttccgcc aagagccgcc tgcagacagc 9120
 ccccgctccc atgccagacc tgaagaatgt caagtccaag atcggtcca ctgagaacct 9180
 gaagcaccag ccgggaggcg ggaaggtgca gataattaat aagaagctgg atcttagcaa 9240
 cgtccagtc aagtgtggct caaaggataa tatcaaacac gtcccgggag gcggcagtgt 9300
 gcaaatagtc taaaaccag ttgacctgag caaggtgacc tccaagtgtg gctcattagg 9360
 caacatccat cataaaccag gaggtggcca ggtggaagta aaatctgaga agcttgactt 9420
 caaggacaga gtccagtcga agattgggtc cctggacaat atcacccacg tccctggcgg 9480
 aggaaataaa aagattgaaa cccacaagct gaccttccgc gagaacgcca aagccaagac 9540
 agaccacggg gcggagatcg tgtacaagtc gccagtgggtg tctggggaca cgtctccacg 9600
 gcatctcagc aatgtctcct ccaccggcag catcgacatg gtagactcgc cccagctcgc 9660
 cacgctagct gacgaggtgt ctgcctccct ggccaagcag ggtttgtgat caggccccctg 9720
 gggcggtcaa taattgtgga gaggagagaa tgagagagtg tggaaaaaa aagaataatg 9780
 acccggtccc cgccctctgc cccagctgc tctcgcagt tgggaattc ggatccagat 9840

cttattaaag cagaacttgt ttattgcagc ttataatggg taaaaataaa gcaatagcat 9900
cacaaatttc aaaaataaag ctttttttc actgcattct agttgtgggt tgtccaaact 9960
catcaatgta ttttatcatg tctggtcgac 9990

<210> 3

<211> 25

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 3
agtaattgaa agagctcaga cgatg 25

<210> 4

<211> 23

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 4
tgtcacctc ttggtcttgg tgc 23

<210> 5

<211> 22

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 5
gtactccacc caagtcgccg tc 22

<210> 6

<211> 23

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 6

gcagcagcat cgaagcttct cag

23

<210> 7

<211> 48

<212> DNA

<213> Artificial

<220>

<223> Missense

<400> 7

gcagcagcat cgaagcttct cagatTTTtac ttccatctgg ccacctcc

48

<210> 8

<211> 20

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 8

ccgccaaagag ccgcctgcag

20

<210> 9

<211> 61

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 9

gcagcagcat cgaagcttct cagattttac ttccacctgg ccacctccta gtttatgatg 60

g 61

<210> 10

<211> 1152

<212> DNA

<213> Homo sapiens

<400> 10

atggctgagc cccgccagga gttcgaagtg atggaagatc acgctgggac gtacggggtg 60

ggggacagga aagatcaggg gggctacacc atgcaccaag accaagaggg tgacacggac 120

gctggcctga aagctgaaga agcaggcatt ggagacaccc ccagcctgga agacgaagct 180

gctggtcacg tgaccaagc tcgcatggtc agtaaaagca aagacgggac tgggaagcgat 240

gacaaaaaag ccaagggggc tgatggtaaa acgaagatcg ccacaccgag gggagcagcc 300

cctccaggcc agaagggcc ggccaacgcc accaggattc cagcaaaaac cccgcccgtt 360

ccaaagacac caccagctc tgggtgaacct ccaaaatcag gggatcgag cggctacagc 420

agccccggct cccagggcac tcccggcagc cgctcccgca ccccgctcct tccaaccca 480

cccaccggg agcccaagaa ggtggcagtg gtcggtactc caccgaagtc gccgtcttcc 540

gccaagagcc gcctgcagac agccccgtg cccatgccag acctgaagaa tgtcaagtec 600

aagatcggct ccaactgagaa cctgaagcac cagccgggag gcgggaaggt gcagataatt 660

aataagaagc tggatcttag caacgtccag tccaagtgtg gctcaaagga taatatcaaa 720

cacgtcccgg gaggcggcag tgtgcaaata gtctacaaac cagttgacct gagcaagggtg 780

acctccaagt gtggctcatt aggcaacatc catcataaac caggagggtg ccagatggaa 840

gtaaaatctg agaagcttga cttcaaggac agagtccagt cgaagattgg gtccctggac 900

aatatcaccc acgtccctgg cggaggaaat aaaaagattg aaaccacaaa gctgaccttc 960

cgcgagaacg ccaaagccaa gacagaccac ggggaggaga tcgtgtacaa gtcgccagtg 1020

gtgtctgggg acacgtctcc acggcatctc agcaatgtct cctccaccgg cagcatcgac 1080

11

atggtagact cgccccagct cgccacgcta gctgacgagg tgtctgcctc cctggccaag 1140
 cagggtttgt ga 1152

<210> 11

<211> 1152

<212> DNA

<213> Homo sapiens

<400> 11

atggctgagc cccgccagga gttcgaagtg atggaagatc acgctgggac gtacgggttg 60
 ggggacagga aagatcaggg gggctacacc atgcaccaag accaagaggg tgacacggac 120
 gctggcctga aagctgaaga agcaggcatt ggagacaccc ccagcctgga agacgaagct 180
 gctggctcag tgacccaagc tcgcatggtc agtaaaagca aagacgggac tggaaagcgat 240
 gacaaaaaag ccaagggggc tgatggtaaa acgaagatcg ccacaccgag gggagcagcc 300
 cctccaggcc agaagggcca ggccaacgcc accaggattc cagcaaaaac cccgcccgtc 360
 ccaaagacac caccagctc tgggtgaacct ccaaaatcag gggatcgag cggctacagc 420
 agccccggct ccccaggcac tcccggcagc cgtccccgca ccccgtccct tccaacccca 480
 cccaccggg agcccaagaa ggtggcagtg gtccgtactc cacccaagtc gccgtcttcc 540
 gccaaagagc gcctgcagac agccccgtg cccatgccag acctgaagaa tgtcaagtcc 600
 aagatcggct cactgagaa cctgaagcac cagccgggag gcgggaaggc gcagataatt 660
 aataagaagc tggatcttag caacgtccag tccaagtgtg gctcaaagga taatatcaaa 720
 cacgtcccgg gaggcggcag tgtgcaaata gtctacaaac cagttgacct gagcaagggtg 780
 acctccaagt gtggctcatt aggcaacatc catcataaac taggagggtg ccagggtgaa 840
 gtaaaatctg agaagcttga cttcaaggac agagtccagt cgaagattgg gtccctggac 900
 aatatcacc acgtccctgg cggaggaaat aaaaagattg aaaccacaa gctgaccttc 960
 cgcgagaacg ccaaagccaa gacagaccac ggggcggaga tcgtgtacaa gtcgccagtg 1020
 gtgtctgggg acacgtctcc acggcatctc agcaatgtct cctccaccgg cagcatcgac 1080
 atggtagact cgccccagct cgccacgcta gctgacgagg tgtctgcctc cctggccaag 1140
 cagggtttgt ga 1152

<210> 12

<211> 19

12

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 12

gcattggaga cacccccag

19

<210> 13

<211> 21

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 13

gcttttactg accatgcgag c

21

<210> 14

<211> 25

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 14

ctggaagacg aagctgctgg tcacg

25

<210> 15

<211> 9990

<212> DNA

<213> Artificial

<220>

<223> PrP/tau transgene construct

<400> 15

```

ggcggccgcg acggatccaa aggcagcaaa aaggcagaga ggggtgatact gggcctggct      60
taagcatttg aaacttcaaa gctcaccccc aattacacac ttcttccaac aagtcacacac      120
ctcctaatta gtgccactct ctgtgggcct acggagagta ttttcattct aactaccaca      180
gttgctgagg aatttaatta aaactacaac cttatcccaa cctagatctt tcagcctttc      240
tgtactacca gagaggggtc atacagcatt gttgtgactc ccattataac ttaaagggaa      300
gctcacacaa agtcagagc cctccatacc ctgcaaatga agaagtacgt tctcaaatcc      360
cttggagcag ggccccactt tggcggcaca aactttaatt tctagacgga acggcatctc      420
tacagaaaga aaagccatgg tatctgcatg ataagtctga aaaggacctg ggcaaatctg      480
cagctgacaa ttccagccat tgctgccact gcgagaaaac cctgctgatg gcagcattgt      540
cagcatcatc tcctccagga acaccggcca tcgagccacg aggacaattg ctgctgctgg      600
agtcaattca tctgccagcc acatcatact ctgggaccgt cactaaccag atccaagcag      660
ccttgaggaa gcatgtcttc tgggtgtgac tgatcccaag ggtgacaac aaggtcctca      720
cagaggcatc ttatgtcaac ctatctacca tgcacggtat aagacacatt ctctctgtg      780
ctgtgtggac actgccatca cagcaacag aaaggaaact cactcactgt gtctgatgtg      840
gtggtgcttg ttaggggagt tctgggcatg tatggcacca tcgcccatga ggactcctgt      900
ggggtcatgc cactctact cctctagaga ccatgaagag atggagaggg aagagcaagc      960
acagatgaca ggctagaact aaagaggagt gtcagggtgag cggacctgaa ctcacggctg     1020
ctcagcctga agtgggtgtg ccatctgcat ctggtatctg gtctgaaggt gcgtggatac     1080
cctctgtgcc cgtccagaag ttctctactg aagacagaaa tgcctgtcca gtcatggaag     1140
aatgattggc agttccact tctcagacca ctgaatgggt cagaacaact actgggtgac     1200
cctaaggtat tcttcagcag atatgtgtga aaaatggaaa gaagatgggt agaaataaac     1260
ggtttttagag gaaaaaaact ctcacaaaga tattataaaa agaaaagagc tttattattg     1320
agcaagcatt caaccagaat gcacaccaca ggcagtctgc taaggagtg tgcagacagg     1380
aggagtgtcg ccctttatgt gagccagtag ataaggatgc tgtgcgtgtt tttagtaact     1440
ggtcttcagc ttgacagcac catttatcac atggtttaac ctaaattcat ctggcgaatg     1500
aggctgtcac gtacttcctg attagcttta tctgaaatga gacaagcttc acatgttcac     1560
ggcaggaggt aatcctgtg cttagagaac agggccatc caagccagge tccttctccc     1620
accaacacgg gtggttgaag agctatctct ccctgggtgtg tgtgtttcag agatggctcc     1680

```

cagggtttttg gtttgggttg aattgggttt tggttttctt actctagccc agactagctt	1740
ggaattctctt ggaagctgc aacggggagc tcagggttcag tgagagatcc tgtctcaaaa	1800
agcagggtga gaagtgattg aggaagacac cccagtgtta acctctgacc tccatatgtg	1860
catgcatgga cagcatgga tacacatata cacacacaca cacacacaca cacacacaca	1920
cacacacaca cacacacaaa accagaaaga atgaacgccc cctctccagc ttgtttacag	1980
tagatacaga gcaactgtaa aacatggggg gtaaaactgaa tgctgagagt aacttagatg	2040
agtaattaag gaaggaagag gaaagaaacc aggaaaccga gagcaagtga ctggaagatc	2100
gttaggcaat ctccacaccc tgctcgttga agttggaatg ctttcttctt ctgcctcttg	2160
aagttcttta gaagtgttag gatttcacaa ttagtctgtg gtggtttcaa tatgcttcac	2220
ccgtggtaag tggcactatt aggaaacgtg tccttggtga aggaagggtg tcaactgcata	2280
ggcgggcttt gaggtgtctt ccagtgtctc agctcctccc agtgcaagag aggcagacac	2340
ctgttgcttg cagaagacag tctcctgctg cctttgaatc aagatgtaga actcaagccc	2400
catgtctgcc tgaacctctg aaactgtaag ccagcccca ttaaatgttt tctttcacia	2460
gagttgcctt ggtcatgggt tctgttcaca gcaataaaac cctaactaag acagtcttaa	2520
atcaatgaaa agacctttaa ttattcattg aacaaacacc attttcttgt atcaagttgg	2580
cagtgtactag taagcaacta tagttctgca ccaggacct ttttgagaa atataccgat	2640
ccaagcatgt tggcatctag attccaaagc caagacacct gccacacct tccatgcctt	2700
gggttctctg caggcatctt ggcttcgggg atgtgtattc caggcaccca ctggaatgca	2760
tggaaacaat taaaatagca tcatagaaga cattgcaatc ctaggagaa actataccaa	2820
aactcagaac tatacctggt taagtgtaga aaagacgaaa ggaataaaac caggaaatatt	2880
ttaaaatatt ttatttgagc tcattgtcat gggatatttg cctgaaagta tgtctgtgta	2940
ccacatgcat ggctggctcc tgcagaggcc aaaagagagc atcagatctc cttagagctg	3000
agtttcagaa gtttgtgagc taccacatgg gtgctggaaa caaaaccag gacatctgga	3060
agagcagcca gtgttcttaa ctactgagcc atcactcagg tcccaccatg aatgtttttc	3120
tttattcttc tctatatttt ctaatgtttt tattggaaat atacaacttt tgccacacat	3180
aacaaatgac caaagaaatg aggtgagagg ggcagctgtt caaatgctgc ctgggaaggc	3240
ttggccagcc ctggcttggc tgccctggc tcagctggcc ctgacttggc tgtcccggtg	3300
ccagctgtca tctactgctt cataataagc tgcactttgg gctgaagggg tggctcagcc	3360
tttaaaggct aggtcataa ccaaagtaag ttgcatttta ttgacttag gttgaagggg	3420
gatctgaaac ttgctgtcaa tgttataaaa cattttatct tcaaatttgg tatagggggc	3480

atagacaaaa ggttctataa accccagaac agcaccactc cctagaaata agcaccata 3540
caagagccta tgggacactt tatagccaaa caaaaagcta tgtttgaaac ttcctttaca 3600
agggcctgag tcccattcat aagggaagga gcccacttc gtaataaac cccactgggtg 3660
acatttgaag gggacacatt caaactgtaa caccatctta tatcatttgc acattaggggt 3720
caaactgtgc cacgttgta tttctaagaa gacagaagtt gtcaagcctg tgctttgagc 3780
cacaagtgtg acaacctact ttcaggcaag tcgctacttc cctaagactc taccaccaata 3840
ggcctgggggt ctggaatgtg ttttaacacag atgcaggett ctgccttagt gcaggcttga 3900
gttctcatgt ccctctctct ttagctttcc gtctcaaggc gcctctcctt agcagaaaaa 3960
atcagaggca taaagcatac atcaggggga agccagagtt ttcagaggga gttttgtgat 4020
ggccttttca gagcattctt gtcaagacta gtttgcctcg ttctctttat taaatgaaag 4080
aaaaataatg cagtgttgca aattagcttt ggtaatggct ccaaccattg tcaggttcac 4140
agtctcatte cgccattcaa aacaacaaac ccaccacact ctctatgcag tgccgtaact 4200
cagaacagcc accaaacagc agaaagaggc tccccgactc ctctcagcct tgccataaac 4260
tcgccggcca catgcttatt ttaaattatt taaattatgt cgtttctccc aacaatgacc 4320
tccaagtgc ttggttgaca ggcttatacc attaagccga ggcttgcata gcaacgataa 4380
ccaggtaggc tattattata accaggtagc tgccgagcta ctggtcggtc cccttttgtc 4440
tctagaaacc tctcaacccc caccacaaaa agcttttatt gccacttcct agtgggtaga 4500
gagcagtcag ccaatagata tttgattctt tgaggaaaaa gctgagtttt gatgtctttt 4560
aatcaagcct ttcagagtcc ctctgtgggg gaggccaggt ggaagcgggg tgggaagctg 4620
gtcccttacc taagctaate tagacacct cccactctc ccctgccctc ttgacagatg 4680
cagtcatcct gatcacaatg agtattctct gaggcaggaa ggcaaggctc tgggaagatgg 4740
tcaatgcctt cattaagaac ccagagtaaa ggtcaagcag acaccagcac cgctgaaatc 4800
taatttctact gtaattgaat catctcagcc aaaggctgta ttttccagcc ctctcgtggc 4860
ctcttcccca acaactgtca acaactgtgt gagcctaccc atgtatgcgc gctcacacac 4920
acacacacac acacacacac acacacacac aggggtggggg gacacaatga ttacacaaga 4980
gtacttaata aacaactata attctcctgg ctcgatagt tcttaccac cctctcctcc 5040
tggtatccga tcttaatact ggatacaaat atttaatcca aaccaatct tgtgtctgtt 5100
aatgatcttc agtgtctcgc cctcagcaag aggacaggat attatgtttt ccctgtgatt 5160
tatgacctct tctgtctcag tatcggcagc aatttattta catggccttg gagtgtgtta 5220
tatgtgtagt atggacatga ggggtcatgt caacctatgt gtggaggcca gaggtcaatg 5280

tcattgtcttc cccaatcaact gccagtggt ccttggtatc caaactcagg tcttcacgct	5340
tggaactga gccagtgccc cagctcctaa cctccccctg ttttaaaaag gtctcattat	5400
gttgcccagg tcagccttga acttgagagt cctctgactg caggctttca cctgtccaag	5460
tcagcaggca tcttgaacaa gaacatcatt tcttttaagc tgtttcaggc tgtgtttggt	5520
gggagctgtt aaatgcagtg ctttttttcc ttggacaca ataaaagaaa aaagtgatta	5580
aatgagttgg gtgtggtggt gcgagtctac aatcctagaa ctcaggagat tgaggagaaa	5640
gcattgctct gagtttgagg tcagcttaaa ttacttagta ggaacaccag gccaaattgg	5700
gctatgggat tgtctccaaa gataaagaaa aaagggaggg agagaaaaga aaaagaaagg	5760
aaagaagggg aaaagaagga atcagcagag aataataaag tcaacatgca atggccaata	5820
tactttctag gcctctaatt cttttatagt ttgtgggaaa atgtcgaaaa tcttcgttac	5880
caatttcttg ttaccaaagt tcaacgatgg cttcctcgct ccgtaggta acctttcatt	5940
ttctcaacta cccattatgt aacgggagca ttgggtactg gatcagtcct ccattaaaga	6000
tgatttttat agttgctgag cgtcgtcagg gagtgctgac actgggggag gtttaaacag	6060
atacaagcat ttaagccagt ccggagcggg gactcatccc cccccacccc ccccccccg	6120
cgagagacgc ggcgcggcca ttggtgagca tcacgccccg cccctcgccc cgcctagttc	6180
ccgcctgccc cgcctcttc cactcccggc tccccgcgt tgtcggatca gcagaccgat	6240
tctgggcgct gcgtcgatc ggtggcagggt aagcgggctg ctgaagccag gccttggcga	6300
gcactcagcc ttccgtcgtc aagctcggct cactgcgcct ctcggggcct tgaggccacg	6360
gggactagga ctgggactgg gactggggct gagtctggct gggaggtgac tgtacacccc	6420
ctgctgcgcg actcctggag gaaccgaatc ccagggcagc caggccggga gccagccttt	6480
ccttcccagc ccagattcac agctcagcat cgctggggat gggggtggca tcttttgact	6540
gtccttggct gttttcttct ctctttgtag tagctacagc gaacataatt ttacctcgtt	6600
attccaccac agtcattact ccttgcaca gtttcattct caacgtcgcc gtgcgccttc	6660
actgcctgt ctaggcgttt tcattgattgt ctattttctt gtactttgaa taccgtgggt	6720
taatagcagt tgcgggtgcg cagaattctc catttcctta agagaaactc ctgggagaat	6780
gggactaaag acgtgcaaat ttaattatat cgcaaacagg aatcaaaatt ttgcattaaa	6840
atgccaaaaca tcttgaaaaa ttaactatc aatgaagaaa aggaactact ttacctacac	6900
acacatccga gagcttcgag gaggcgaagg aaatagaaa ctaagggatg atttggttg	6960
tatttgaatc tgacacaagc ttccatatt atttatagea gggactaaac gatgagtcac	7020
tttctgaata agatgcaaat taaagcaagt ttgtttgttg tctttacac tattaaatag	7080

acagagacaa tggcaacagc aaccttaacc tagaggttgc ctgaaagtgt cagggtttggg 7140
 aacaagtggc cctgcttaag ggctagaaag attgctttac aaccaacaat catgacttga 7200
 cattgcctgg ggttcctttt gtcctttcct tttttaaaag actagtgttt atcttatgtg 7260
 catgagtgtt ttgcatccac attcgccctgt atacacacct ggttctgtgg aggtcaggag 7320
 aggggtgctgg atgccctggc actagagcct tggatggtta tgtgagcccc tggcacaggg 7380
 gagctcagaa ccaaattccag gtccctctgga agagcaacca gagctcttaa aactttctaag 7440
 tatccctcca tcccttttcc atcatatttg gaaaggagaa aactgctacc catgcctggc 7500
 atttatttca gagattaact gtctgtgtaa aacttgacat tgaaagtga ctattctgtt 7560
 tcccatcat acttagttga gactactgta agtcagttag ggcttttttt gtttgggtcc 7620
 ttggttagtt tggagtgtgt ttgtgagctc attaacaggc tttcaatatg tagctggaat 7680
 ttgctgtgta gaccagacag gcctcaaatt tgtggcaatc ctccctgcat ctcccagaa 7740
 tgccttggtg caggcataaa ccaccgtgcc cagcagtaaa acaatctggt gaggtattat 7800
 tagtcgtgtg ctgtgacca gaaacccac tcctggcaat ttactgggaa ggaacaaaca 7860
 aagggtctagg ggagccatat ggctgcagt tagagaaaat tagatccaac tgaaaaatca 7920
 acctaaagggt gtaaaagcca agcagttaag aaactgacaa gctcatgatg gaagccgagg 7980
 ccatcgtgaa cactcttcat tttaggcccc acgtatcact ggggacaact gagagtcaaa 8040
 gtacaggtaa ggagaccaag gcttttcagg actcaggctg tctcagtga aagcccagaa 8100
 gagcagtaat tgaaagagct cagacgatgt gtctgatctc ctctgtttgt ttgttctgt 8160
 attatttcca ctaacttatt tgggaggaaa aaaaacagtt cacaggcttc tttcttgaa 8220
 atactgggga ttgctgggat cgaaccagag gataggtttt tagtttctaa aataacatag 8280
 atcatgcctt gtttgccttt tggaatatgt ttgcgtgcc cttattttca tgttcaaata 8340
 ctgctccatt ttgcgtgact ctttagtatt gggttgatga ttgcatatt agattagatt 8400
 gtatttcagt tctcagactt atttatcaat tctagttttc tctttttgtt gttttaaagg 8460
 actcctgagt atatttcaga actgaacct ttcaaccag ctgaagcatt ctgccttcct 8520
 agtggtagct cgactatcag gtgaactttg aaccaggatg gctgagcccc gccaggagtt 8580
 cgaagtgatg gaagatcacg ctgggacgta cgggttgggg gacaggaaag atcagggggg 8640
 ctacaccatg caccaagacc aagaggggtga cacggacgct ggcttgaaag ctgaagaagc 8700
 aggcattgga gacaccccca gcctggaaga cgaagctgct ggtcacgtga cccaagctcg 8760
 catggtcagt aaaagcaaag acgggactgg aagcgatgac aaaaaagcca agggggctga 8820
 tggtaaacg aagatcgcca caccgcgggg agcagccct ccaggccaga agggccaggc 8880

caacgccacc aggattccag caaaaacccc gcccgctcca aagacaccac ccagctctgg 8940
tgaacctcca aaatcagggg atcgagcggt ctacagcagc cccggctccc caggcaactcc 9000
cggcagccgc tcccgcaccc cgctcccttc aacccccccc acccgggagc ccaagaaggt 9060
ggcagtggtc cgtactccac ccaagtcgcc gtcttccgcc aagagccgcc tgcagacagc 9120
ccccgtgccc atgccagacc tgaagaatgt caagtccaag atcggctcca ctgagaacct 9180
gaagcaccag ccgggaggcg ggaagggtgca gataattaat aagaagctgg atcttagcaa 9240
cgtccagtc aagtgtggct caaaggataa tatcaaacac gtcccgggag gcggcagtg 9300
gcaaatagtc taaaaaccag ttgacctgag caagggtgacc tccaagtgtg gctcattagg 9360
caacatccat cataaaccag gaggtggcca ggtggaagta aaatctgaga agcttgactt 9420
caaggacaga gtccagtcga agattgggtc cctggacaat atcacccacg tccctggcgg 9480
aggaaataaa aagattgaaa ccacaagct gacctccgc gagaacgcca aagccaagac 9540
agaccacggg gcggagatcg tgtacaagtc gccagtgggt tctggggaca cgtctccacg 9600
gcatctcagc aatgtctcct ccaccggcag catcgacatg gtagactcgc ccagctcgc 9660
cacgctagct gacgaggtgt ctgcctccct ggccaagcag ggtttgtgat caggccccctg 9720
gggcggtcaa taattgtgga gaggagagaa tgagagagtg tggaaaaaaa aagaataatg 9780
accgggcccc cgccctctgc cccagctgc tctcgcagt tcgggaattc ggatccagat 9840
cttattaaag cagaacttgt ttattgcagc ttataatggt taaaaataaa gcaatagcat 9900
caciaatttc aaaaataaag ctttttttc actgcattct agttgtggtt tgtccaaact 9960
catcaatgta tcttatcatg tctggtcgac 9990